

Microrna Regulation of Nodule Zone-Specific Gene Expression in Soybean

Md Ehsanul Haque^{1*}, Most Shanaj Parvin²

¹South Dakota State University, Brookings, SD, USA

²Bangladesh Agricultural Research Institute, Joydebpur, Gazipur, Bangladesh

Article Info

Received: June 05, 2021

Accepted: June 14, 2021

Published: June 21, 2021

***Corresponding author:** Md Ehsanul Haque, South Dakota State University, Brookings, SD, USA.

Citation: Md Ehsanul Haque, Most Shanaj Parvin. (2021) "Microrna Regulation of Nodule Zone-Specific Gene Expression in Soybean.", Journal of Agricultural Research Pesticides and Biofertilizers, 2(1); DOI:<http://doi.org/06.2021/1.1028>.

Copyright: © 2021 Md Ehsanul Haque. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract:

Nitrogen is a paramount important essential element for all living organisms. It has been found to be a crucial structural component of proteins, nucleic acids, enzymes and other cellular constituents which are inevitable for all forms of life. In the atmosphere, the percentage of nitrogen is very high (N₂, 78%) compared to other inorganic gases. However, most organisms have practically no direct access to this nitrogen. While plants can not directly uptake nitrogen from atmosphere, they are capable of assimilating other forms of nitrogen, for example ammonium (NH₄⁺) and nitrate (NO₃⁻). For agricultural crop production, artificial fixation of nitrogen is heavily utilized, and it is an expensive process that requires high temperatures (at least 400 °C) and pressures (around 200 atm). It has been conspicuously demonstrated that indiscriminate use of fertilizer hampers soil physical, chemical and micro biological properties and also a potential risk to environment e.g. water quality. Besides, chemically manufactured fertilizers are depleted from soils in various ways, for instance; denitrifying bacteria, volatilization, and leaching. Consequently, it results relatively poor availability of nitrogen to get into plants. On the flipside, only 1-2% of the nitrogen fixation in the world occurs through the natural process of lightening. Notably, microbial fixation is well characterized in diazotrophs for example; Rhizobia and Frankia, and blue-green algae. Against the backdrop, we are accentuated on an environmentally friendly and the most sustainable approach to increase productivity for legume and non-legume crops. Till today, the term biological nitrogen fixation (BNF) has received much attention as a sustainable alternative; this process facilitates atmospheric nitrogen to convert into ammonia by rhizobia in specialized plant organs termed "root nodules". This review article seeks to better understand plant mechanisms involved in the development of root nodules in soybean.

Introduction:

Soybean (*Glycine max*) is one of the most important oil crops and a source of animal feed protein in the world. It has a salient feature to fix atmospheric nitrogen through symbioses with compatible rhizobia that yields to determinate type nodule (Oldroyd, Murray et al. 2011). Biological nitrogen fixation in soybean nodules reduces the use of chemical nitrogen fertilizers resulting in cost-savings to producers and minimizes environmental damage due to nitrogen run-off. A better understanding of how nodules form and function is important for selection or generation of soybean genotypes with better nitrogen fixation capacity. Soybean nodules originate from root cortex via de novo cell differentiation (Oldroyd 2013). Consequently, two major nodule development zones are formed for instance; the nodule primordium (Npr) in the middle and it is encircled by nodule parenchyma (Npa). At later time point, the Npr gives rise to N-fixation zone and the Npa holds vascular bundles. It is not clear what early signaling pathways driving the conspicuous development of the nodule zones. My research is aimed at filling this knowledge gap by illustrating the molecular signatures that paves the way to cellular differentiation in root nodule development in soybean. Based on initial evidence obtained by the Subramanian lab, we hypothesize that microRNAs (miRNAs) play important regulatory roles in spatio-temporal expression of their target genes during nodule developmental in soybean. For instance, the regulation of auxin sensitivity by miR160 has been found to be crucial for formation of nodule primordia and vasculature in the parenchyma (Marie Turner 2013). Against this backdrop, this review article focused on nuclear and cytoplasmic transcriptome as well



miRNA profiles of parenchyma and primordial tissues and determine the relative abundance and differentially expressed mRNAs and regulatory role of miRNAs in cell differentiation and nodule development.

Root nodule a sustainable alternative to fix atmospheric nitrogen:

Atmospheric nitrogen percentage is very high (N₂, 78%) compared to other inorganic gases (Mary Elvira 1932). However, most of the organisms have practically no direct access to this nitrogen. Nevertheless, plants can not directly uptake nitrogen from atmosphere, but they are capable of assimilating only very specific forms of nitrogen, for example ammonium (NH₄⁺) and nitrate (NO₃⁻) (Bytnerowicz and Fenn 1996, Peter M. Vitousek 1997) (Sponseller, Gundale et al. 2016). Virtually, nitrogen has been found to be a crucial structural component of proteins, nucleic acids, enzymes, and other cellular constituents which are inevitable for all forms of life (O'Brien, Vega et al. 2016). For agricultural crop production, artificial fixation of nitrogen is heavily utilized. It is an expensive process that requires high temperatures (approx. 400 °C) and pressures (approx. 200 atm) (Witschi 2000). It has been conspicuously demonstrated that indiscriminate use of N fertilizer hampers the diversity of the bacterial community and decreases soil C and N concentrations (Verzeaux, Alahmad et al. 2016). Notably, it has been demonstrated as a potential risk to environment e.g. water quality (Zhao, Sha et al. 2016) (Sponseller, Gundale et al. 2016). Besides, chemically manufactured fertilizers are depleted from soils in various ways, for instance; denitrifying bacteria, volatilization, and leaching (Johnson 1996, Peter M. Vitousek 1997). Consequently, it results relatively poor availability of nitrogen to get into plants. On the flipside, over 90 % of the nitrogen fixation in the world occurs through the natural process of lightning and microorganisms. Furthermore, microbial fixation is well characterized in diazotrophs for example; Rhizobia and Frankia, and blue-green algae (Cheng 2008). It has been demonstrated that Bradyrhizobium strains substantially escalated soybean grain yield, and protein content up to 57% and 26%, respectively (Zimmer, Messmer et al. 2016). Against the backdrop, we are accentuated on an environmentally friendly and a sustainable approach to increase the productivity for legume and non-legume crops. Literature mining depicted that biological nitrogen fixation in soybean nodules reduces the use of chemical nitrogen fertilizers resulting in cost-savings to producers and minimizes environmental damage due to nitrogen run-off.

Rhizobia infection leads to the root nodule development:

In the natural environment, plants are continuously confronted with pathogenic and symbiotic microbes. Symbioses involves mutual exchange of diffusible signal molecules, first endophytic bacteria (rhizobia) are attracted by the plant root exudates flavonoids which are perceived and triggered the bacterial nodulation (nod) genes. Consequently, the bacteria synthesize specific lipochito-oligosaccharides, called nodulation (Nod) factors. This signal is perceived by the LysM receptor like kinase of host plant, it induces the root hair curling, and bacteria get access into the host epidermis through infection threads (ITs) and initiate cell division within the root cortex, leading to the progression of the root nodule meristem. In later stages of the interaction, bacteria are released from the infection threads into the

plant cells, surrounded by membrane of plant origin. These bacteria multiply within the host cells and differentiate into the nitrogen fixing bacteroids (Udvardi and Day 1997) (Oldroyd 2013). Till now, integration of genetic and genomic approaches has revealed twenty-six genes to be involved in nodule development of Medicago truncatula and Lotus japonicum (Kouchi, Imaizumi-Anraku et al. 2010). In addition, deep sequencing of the Medicago truncatula root transcriptome has uncovered thousands of genes to be induced during Nod factor signaling and its resulting ethylene (ET) biosynthesis throughout the multiple development stages of indeterminate nodule (Larrainzar, Riely et al. 2015). Albeit the molecular mechanism of such regulation is not well understood. There has been a large-scale transcriptome analysis of *B. japonicum*-inoculated and mock-inoculated soybean root hairs. It has showed that a total of 1,973 soybean genes differentially expressed during root hair infection, particularly NFR5 and NIN genes (Libault, Farmer et al. 2010). Nevertheless, the signaling mechanisms directing the cellular differentiation of nodule are not known.

Soybean root nodule organogenesis:

Soybean (*Glycine max*) has a genome size of 1.1 to 1.5 Gb, it is partially diploidized tetraploid. It is one of the most important oil crops and a source of animal feed protein in the world (soybase.org/sb_about.php). It has a salient feature to fix atmospheric nitrogen through symbioses with compatible rhizobia that yields to determinate type nodule (Udvardi and Day 1997) (Oldroyd, Murray et al. 2011). Notwithstanding of the economic and environmental importance, there has been very few studies about quantitative trait loci (QTL) that controlling BNF traits, for instance nodule number, ration of nodule dry weight with nodule number, and shoot dry weight (SDW). It has been reported via composite interval mapping that approximately six QTLs bears very small effect on BNF traits (Santos, Geraldi et al. 2013). Besides, it has been demonstrated in earlier studies that nodules originate from root cortex via de novo cell differentiation into two different cell types, parenchymal and primordium (Celine Charon 1997) (Oldroyd & Downie 2008; Oldroyd 2013). In addition, early nodulin genes in legume for instance; Enod 40 gene reported to be expressed in root pericycle during the rhizobia infection and later it occupied in the dividing cortical cells (H. Kouchi and S. Hata 1993). Among the two major nodule development zones, the nodule primordium (Npr) in the middle which is encircled by nodule parenchyma (Npa). At later time point, the Npr gives rise to N-fixation zone and the Npa holds vascular bundles. Lately, a β - expansin gene, GmEXPB2 fused with GUS reporter gene which was observed to be preferentially expressed in nodule vascular trace and nodule vascular bundles. It indicated that GmEXPB2 might be crucial for nodule organogenesis. Over expression of GmEXPB2 contrast to suppressed GmEXPB2 transgenic lines found to be escalated nodule number, nodule mass and nitrogenase activity. It further suggested that GmEXPB2 might have influenced over root architecture, nodule formation and development, and profoundly yielding to biological N₂ fixation (Li, Zhao et al. 2015). Even though, it is not clear what early signaling pathways driving the conspicuous development of the nodule zones. Against the backdrop, to understand the regulation of auxin sensitivity by miR160 which is believed to be crucial for the formation of nodule primordia (Marie Turner 2013).




Figure 1a: Illustrating the progression of root nodule development through Rhizobial bacterial infection in the plant root leading to the determinate nodule (Oldroyd 2013).




Figure 1b: Nodule development zones A. Nodule primordial zone (Enod 40 gene) in the middle B. surrounding parenchyma (Enod 2 gene), differentiated from cortex (collected from Sen Subramanian lab).

Regulatory small RNAs biogenesis and its molecular functions:

Regulatory small RNAs are ranged between 20 to 24 nucleotides which are ubiquitous elements of endogenous plant transcriptomics, a common response to exogenous viral infections and introduced double-stranded RNA (Axtell 2013). Three core enzymes families, for instance; RNAdependent RNA polymerase (RDR), Dicer like (DCL), and Argonaute (AGO) proteins paves the way of small RNA biogenesis and function in plants. Firstly, ribonuclease type III or DICERLIKE1 involves in the yield of a fold-back precursor RNA or primary miRNA (pri-miRNA) transcripts using an RNA templates in the nuclei. Later, the resulting miRNA-miRNA duplex which is originated in nucleus then translocated into cytoplasm. The guided miRNA molecule is incorporated into ARGONAUTE (AGO) to form an active RISC complex to specific target RNAs that are complementary to the miRNA, and this process eventually follows up mRNA cleavage, represses the translation of the mRNAs or Chromatin modification. This phenomenon accentuated as an inhibition or silencing of the gene expression, which play a crucial role in the developmental process in plant and animal (Chapman and Carrington 2007) (Axtell 2013).




Figure 2: Regulation of gene expression events via RISC complex




Figure 3: Gene expression events occurring in typical plant cell

It has been found in several studies that most plant miRNAs are non-coding RNA, and small 21-24 nucleotide long (Cuperus, Fahlgren et al. 2011). It requires DCL1-clade DCL for their biogenesis and AGO1-clade AGO for their function (Wu, Zhou et al. 2010, Manavella, Koenig et al. 2012).

In rice (*Oryza sativa*), DCL3 has been reported in the biogenesis of 24nt long miRNA that incorporated in AGO4 to regulate the target gene expression primarily through mRNA cleavage (Wu, Zhou et al. 2010). Argonaute proteins (AGO) form RNA inducing silencing complexes (RISC) with small RNAs which is known as post-transcriptional gene silencing. It has typically four domains, for instance: N-terminal, PAZ, MID and PIWI domains. The MID-PIWI lobes are belongs to the C-terminus. It has been studied that MID-domains contains the specificity loop to recognize and bind to the 5'-phosphate of smRNAs. The PIWI domains contained the catalytic active site D-E-D-H/D. PAZ domain anchored the 2-nt overhang at the 3' end of miRNAs. The N-terminal domain involved in the separation of miRNA-miRNA duplex and the slicer activity of the mRNA (Song, Smith et al. 2004). There has been an expansion and duplications of AGO



family members during plant evolution (Singh, Gase et al. 2015). soybean mosaic virus and among them

The functional diversification of AGOs is indicating 71 genes projected to induce in defense response (Hui Chen 2016). sRNA directed regulatory pathways. The binding preference of These findings suggested the

AGO and sRNA is mainly assigned by the sequence of sRNA. In regulatory role miRNA that governed the target gene expression

Arabidopsis, 10 AGO have been extensively studied (Liu et al. during viral infection.

2014). It has been demonstrated that AtAGO10 like AtAGO1, it Furthermore, the regulatory role of microRNAs (miRNAs) during

recognized distinct structural features in miR165/miR166 duplex Soybean- Bradyrhizobium

than involved by AtGO1. AtAGO10 found to regulate shoot apical japonicum mutualistic association was studied first by

meristem by decoying miR165/miR166 and subsequent repression Subramanian et al. 2008. They sequenced

of homeodomain-leucine zipper (HD-ZIP) gene expression (Zhu, approximately 350000 small RNAs of soybean root sample which

Hu et al. 2011). Notably, 22 AGO proteins have been reported in were inoculated with B.

Soybean (*Glycine max*). It has been found that genome duplication japonicum. It helps to detect 20 conserved miRNAs loci based on

in Soybean resulted such a proliferation of AGOs. For example: its the similarity to miRNAs in

genome encodes two copies of AGO1, AGO2, AGO5, AGO4/9, another plant species. In addition, 35 novel miRNAs were

AGO6 and AGO7 (Xiang Liu 2014). However, the molecular identified based on potential hairpin

function of the plant AGO genes yet not very clear. forming precursors in Soybean EST as well as shotgun genomic

There are several miRNA families that are conserved across the sequences (Subramanian, Fu et

vast evolutionary distances from al. 2008). These findings advocated the potential role of miRNAs

flowering plants to mosses (Cuperus, Fahlgren et al. 2011). It has in the regulation of legume rhizobium symbiosis.

been observed in another study that miRNA, and its target pairing found to be stable for a

prolonged periods of plant evolution. In another study, 120 hairpin-forming precursor genes have been

On the flip side, another group demonstrated that conserved plant identified in soybean by Turner

miRNAs and their targets are to somehow flexible. For instance; et al. In addition, they reported three novel miRNAs for instance;

miR159 is a highly conserved miRNA that targets not only a miR160, miR164 and miR393

subset of MYB mRNAs but also observed to target a non MYB found to be involved in auxin signaling (Turner, Yu et al. 2012).

mRNA, SGN-U567133 (Buxdorf, Hendelman et al. 2010). A Moreover, the plant hormone

mutant tomato transgenic line (miR159-resistant line) showed auxin is thought to have a pivotal role in nodule organogenesis in

higher level of the SGN-U567133 transcript and exhibited defects determinate and indeterminate

in leaf and flower development. This result suggests that miR159 type of nodule. It indicates a redundancy and diversity of miRNAs

involves in a post-transcriptional regulation. Additionally, it is family members that governs

found to be crucial for the normal tomato development. the formation of root nodule.

Recently, the identification of miRNAs in the regulation of It has been illustrated that auxin receptor gene family hushed by

photoperiodic pathways in soybean overexpressed microRNA393.

have been reported through high throughput sequencing and qRT- These plant roots found to be hypersensitive to auxin and yielded

PCR. Six libraries were normal nodule. This observation

constructed using Illumina Solexa, for instance; 0, 8, and 16 h advocated that only minimal/reduced auxin signaling is required

under short day treatment, similar for determinate nodule

time points considered for the long the long day treatment. A total development. Likewise, overexpressed microRNA160 hushed a

of 163 miRNAs families were set of repressor auxin response

reported which covered 318 plant miRNAs, and unclassified 81 transcription factor. These plant roots were hypersensitive to auxin

novel predicted miRNAs. As and observed not to be reluctant

expected, significant differences in abundance between short day in epidermal responses to rhizobia. Notably, it yielded to lower

and long day treatment was sized nodule primordium (Marie

observed (Wenbin Li 2015). These findings provided evidence of Turner 2013). This observation indicated that auxin

miRNA in the regulation of hypersensitivity inhibits nodule organogenesis

flowering time that ultimately affects the seed yield and quality of Organ specific expression of profile of miRNA and the potential

soybean. targets were also studied. Two

The complex regulatory network of miRNA-mRNA interactions genes (*Glyma10g10240* and *Glyma17g05920*) which were the

during viral infection has been highly expressed in soybean nodule. Likewise, three potential

revealed via small RNA seq (sRNA), degradome seq, and genome- targets of *gma-new-miR13587*

wide transcriptome analysis. demonstrated to be highly expressed in the nodules than in the

There has been a total of 253 soybean miRNAs found to be two- roots. As expected, *gma-newmiR13587* found to be poorly

fold abundance compared with expressed in the nodules than in the roots (Turner, Yu et al. 2012).

mock-inoculated control demonstrated through sRNA seq There was an inverse expression pattern observed in between roots

analysis. Among them 105 miRNAs and nodules.

were identified as potential targets of 125 transcripts that has been Li et al., studied the transgene expression of three novel miRNAs

validated by degradome seq namely, miR482, miR1512, and

analyses. In addition, 2679 genes were detected via genome wide miR1515 in Soybean. They noticed a significant increase of nodule

transcriptomic analysis. These numbers while root length and

genes have been differentially expressed during infection of later root density were normal in all tested miRNA lines. As

expected, there were differential



expression of these miRNAs in supernodulating and nonnodulating soybean mutants. They reported that 6 novel miRNAs decayed 22 predicted target genes. And it was estimated via real time polymerase chain reaction and qRT-PCR (Li, Deng et al. 2010). It advocates that miRNAs have the signatory roles in soybean nodule development. Sequencing of small RNAs and Parallel analysis of RNA ends (PARE) libraries revealed to identify 284 nodule miRNAs, more than 500 target genes, and including 178 novel soybean miRNAs. It has been reported that ENOD93 only found to be expressed in nodule tissue not in other plant parts of Soybean. Ectopic expression of miR393j-3p and RNAi silencing approach to ENOD93 expression showed a significant reduction in nodule formation (Zhe Yan 2015). Therefore, this study showed a list of miRNAs and their potential target of nodulation genes. In the model legume (*Medicago truncatula*), 25 conserved miRNA families and 100 novel miRNA reads were detected by high-throughput sequencing. The expression of MtHAP2-1 (encodes a CCAAT binding transcription factor) to meristematic zones was restricted by miR169a which is found to be critical for the development of indeterminate type of nodule (Combiér, Frugier et al. 2006). In another study, HDZ1 transcripts were inhibited by overexpression of miR166, it dropped the number of symbiotic nodule and lateral root (Boualem, Laporte et al. 2008). To get insights into key genes of nodule zones, transcript profiles of specific cells/tissues were investigated at different time points from indeterminate nodules of *M. truncatula* using laser capture micro dissection. It has been demonstrated from the comprehensive gene expression map that selected genes enriched in different cell/tissue types (Limpens, Moling et al. 2013). These findings indicated that organ specific gene expression could be controlled by the presence or absence of miRNAs. Recently, *Agrobacterium rhizogenes* mediated hairy root transformation has been applied as tool for exploring cell type specific gene expression in tomato. Cell type or tissue specific promoter introduced into INTACT and TRAP constructs via gateway cloning technology to develop binary vectors. INTACT method used to capture biotin tagged nuclei from specific cell types and TRAP method used for profiling of mRNAs or foot printing of individual ribosomes (Ron 2014). TRAP methodology is not required tissue fixation or single cell suspension. It has been successfully used to date in organisms ranging from *D. melanogaster* to mice and human cultured cells. Multiple ribosomes or Polyribosomes (polysomes) are engaged in translation on a single mRNA. To evaluate the translation state of an mRNA, ribosomal subunits, ribosomes, and polysomes can be isolated from detergent-treated cell extracts (Heiman, Kulicke et

al. 2014). In this study, we would perform polysome isolation deploying gene cassettes ENOD40p:HF-GFP-RPL18 for primordial tissues, and ENOD2p:HF-GFP-RPL18 for parenchymal tissues in *Glycine max* root nodules that express an epitope tagged version of ribosomal protein L18. Over the last one decade, there has been several microarrays-based studies which characterized transcriptional variations deployed in nodule formation. It has been embedded with couple of short comings, for instance; relative late time points study, incomplete representation of plant genes, discrimination of close paralogs, and reduced sensitivity. Lately, next generation sequencing technology have widened the horizon of transcription analyses in different legume species to detect symbiosis induced changes in late nodule developmental stages. Against this backdrop, we are accentuated to reveal early transcriptional changes induced in determinate type of soybean nodule by *Bradyrhizobium japonicum*. In determinate type of nodule, two major nodule development zones are formed for instance, the nodule primordium (Npr) in the middle and it is encircled by nodule parenchyma (Npa). At later time point, the Npr converted to N-fixation zone and the Npa contained vascular bundles. Of these facts, it is not clear what early signaling pathways driving the conspicuous development of the nodule zones. In this context, mechanisms regulate the distinct gene expression profiles in Npr and Npa cell types has not understood clearly. The proposed research study is aimed at filling this knowledge gap by illustrating the molecular signatures that paves the way to cellular differentiation in root nodule development in soybean considering four different time points (5 dai, 7 dai, 10 dai & 14 dai). The hypothesis is microRNAs (miRNAs) play important regulatory roles in spatio-temporal expression of their target genes during nodule developmental in soybean. For example, a gradient of microRNA localization between nodule primordium and parenchyma cells could result in distinct differentiation of these cell types. To test this hypothesis, one has to obtain both cell type-specific miRNA and transcriptome (miRNA target) profiles. Since, the majority of miRNA regulation occurs in the cytoplasm, we reasoned that comparison of nuclear and ribosomal transcriptome profiles would reveal genes whose expression is potentially regulated by post transcriptional mechanisms such as miRNA cleavage. Combining this information with cell type-specific miRNA profiles, and to test the above hypothesis and identify key miRNA-target pairs important for nodule cell differentiation. The use of translating ribosome affinity purification (TRAP) of nodule zone cells, namely from parenchyma and primordial tissues, to obtain cytoplasmic transcriptomes data.

Techniques to determine cell type specific expression profiles: TRAP methods:

TRAP is termed translating ribosome affinity purification, combines cell-type-specific transgene expression with affinity purification of translating ribosomes. It supersedes the need for tissue fixation, and facilitates to study the cell type-specific mRNA profiles of any genetically defined cell type. It has been successfully used to date in organisms ranging from *D. melanogaster* to mice, and human cultured cells. Multiple ribosomes or Polyribosomes (polysomes) are engaged in



translation on a single mRNA. To evaluate the translation state of an mRNA, ribosomal subunits, ribosomes, and polysomes can be isolated from detergent-treated cell extracts.

In this study, the polysome isolation using gene cassettes ENOD40p:HF-GFP-RPL18 for primordial tissues, and ENOD2p:HF-GFP-RPL18 for parenchymal tissues in Glycine max root nodules that express an epitope tagged version of ribosomal protein L18 RPL18(Heiman, Kulicke et al. 2014, Ron 2014).

Relative abundance and differentially expressed mRNAs profile in two different tissue specific zones would help to understand the effect of regulatory role of miRNAs in cell differentiation and nodule development.

References:

- Axtell, M. J. (2013). "Classification and comparison of small RNAs from plants." *Annu Rev Plant Biol* 64: 137-159.
- Boualem, A., et al. (2008). "MicroRNA166 controls root and nodule development in *Medicago truncatula*." *Plant J* 54(5): 876-887.
- Buxdorf, K., et al. (2010). "Identification and characterization of a novel miR159 target not related to MYB in tomato." *Planta* 232(5): 1009-1022.
- Celine Charon, C. J., Eva Kondorosi, Adam Kondorosi and Martin Crespi (1997). "enod40 induces dedifferentiation and division of root cortical cells in legumes." *Proc. Natl Acad. Sci. USA*. 94: 8901-8906.
- Chapman, E. J. and J. C. Carrington (2007). "Specialization and evolution of endogenous smallRNA pathways." *Nat Rev Genet* 8(11): 884-896.
- Cheng, Q. (2008). "Perspectives in biological nitrogen fixation research." *J Integr Plant Biol* 50(7): 786-798.
- Combiér, J. P., et al. (2006). "MtHAP2-1 is a key transcriptional regulator of symbiotic nodule development regulated by microRNA169 in *Medicago truncatula*." *Genes Dev* 20(22): 3084-3088.
- Cuperus, J. T., et al. (2011). "Evolution and functional diversification of MIRNA genes." *PlantCell* 23(2): 431-442.
- Hiroshi Kouchi¹, K.-i. T., Rollando B. So², Jagdish K. Ladha² and Pallavolu M. Reddy² (1999). "Rice ENOD40: isolation and expression analysis in rice and transgenic soybean root nodules." *The Plant Journal* 18(2): 121-129.
- Johnson, D. S. O. a. G. V. (1996). "Fertilizer Nutrient Leaching and Nutrient Mobility: A Simple Laboratory Exercise." *J. Nat. Resour. L. ife Sci. Educ* 25(2): 128-131.
- Kouchi, H. and Hata, S. (1993) Isolation and characterization of novel nodulin cDNAs representing genes expressed at early stages of soybean nodule development. *Mol. Gen. Genet.* 238, 106–119.
- Li, H., et al. (2010). "Misexpression of miR482, miR1512, and miR1515 increases soybean nodulation." *Plant Physiol* 153(4): 1759-1770.
- Manavella, P. A., et al. (2012). "Plant secondary siRNA production determined by microRNA duplex structure." *Proc Natl Acad Sci U S A* 109(7): 2461-2466.
- Marie Turner, e. a. (2013). "Ectopic Expression of miR160 Results in Auxin Hypersensitivity, Cytokinin Hyposensitivity, and Inhibition of Symbiotic Nodule Development in Soybean." *Plant Physiology* 162(2013): 2042–2055.
- Oldroyd GE, Downie JA. (2008). "Coordinating nodule morphogenesis with rhizobial infection in legume. *Annual Review of Plant Biology* 59:519-546.
- Singh, R. K., et al. (2015). "Molecular evolution and diversification of the Argonaute family of proteins in plants." *BMC Plant Biol* 15: 23.
- Song, J. J., et al. (2004). "Crystal structure of Argonaute and its implications for RISC slicer activity." *Science* 305(5689): 1434-1437.
- Sponseller, R. A., et al. (2016). "Nitrogen dynamics in managed boreal forests: Recent advances and future research directions." *Ambio* 45 Suppl 2: 175-187.
- Subramanian, S., et al. (2008). "Novel and nodulation-regulated microRNAs in soybean roots." *BMC Genomics* 9: 160.
- Turner, M., et al. (2012). "Genome organization and characteristics of soybean microRNAs." *BMC Genomics* 13: 169.
- Udvardi and Day (1997). "Metabolite transport across symbiotic membranes of legume nodules." *Annual Review of Plant Physiology and Plant Molecular Biology* 48: 493-523.
- Weeks, Marry Elvira (1932). "The discovery of the elements. IV. Three impotent gases". *Journal of Chemical Education*. 9 (2): 215
- Wu, L., et al. (2010). "DNA methylation mediated by a microRNA pathway." *Mol Cell* 38(3): 465-475.
- Xiang Liu, T. L., Yongchao Dou, Bin Yu, and Chi Zhang (2014). "Identification of RNA silencing components in soybean and sorghum." *BMC Bioinform* 15: 4.
- Zhe Yan, M. S. H., Siwaret Arikat, Oswaldo Valdes-Lopez, Jixian Zhai, Jun Wang¹, Marc Libault¹, Tieming Ji, Lijuan Qiu, Blake C. Meyers and Gary Stacey (2015). "Identification of microRNAs and their mRNA targets during soybean nodule development: functional analysis of the role of miR393j-3p in soybean nodulation." *New Phytologist* 207: 748–759.
- Zhu, H., et al. (2011). "Arabidopsis Argonaute10 specifically sequesters miR166/165 to regulate shoot apical meristem development." *Cell* 145(2): 242-256.