

Hollow Viscus Injury from Blunt Abdominal Trauma in Children: Experience in a Developing Country

Kevin Emeka Chukwubuike

Department of Surgery, Enugu State University Teaching Hospital, Enugu, Nigeria.

Article Info

Received: October 11, 2021

Accepted: October 29, 2021

Published: November 02, 2021

***Corresponding author:** Kevin Emeka Chukwubuike, Department of Surgery, Enugu State University Teaching Hospital, Enugu, Nigeria.

Citation: Kevin Emeka Chukwubuike (2021) "Hollow Viscus Injury from Blunt Abdominal Trauma in Children: Experience in a Developing Country". *Clinical Research and Clinical Case Reports*, 2(3); DOI: <http://doi.org/04.2021/1.1037>.

Copyright: © 2021 Kevin Emeka Chukwubuike. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Objective: To evaluate our experience in management of children who sustained hollow viscus injury following blunt abdominal trauma.

Methodology: This was a retrospective study of children aged 15 years and below who were managed for hollow viscus injury between January 2009 and December 2018 at the pediatric surgery unit of Enugu State University Teaching Hospital (ESUTH) in Enugu, Nigeria.

Results: A total of 583 cases of blunt abdominal traumas were recorded during the study period, out of which 21 (3.6%) cases had hollow viscus injury. The ages of the patients ranged from 8 to 14 years, with a median of 10 years. Road traffic accident and abdominal pain were the most common mechanism of injury and presenting symptom respectively. Six (28.6%) patients were in a clinical state of shock at presentation. Small intestinal was the most injured hollow viscus and the spleen was the most associated injured solid organ. Surgical site infection was the most common post-operative complication and 1 (4.8%) patient expired. **Conclusion:** Our experience revealed that although hollow viscus injury is uncommon in children, the small intestine is the most injured. Hollow viscus injury can be associated with significant morbidity and mortality.

Key words: Abdominal trauma, experience, hollow viscus, tertiary hospital.

1. Introduction

Trauma to the abdomen could be blunt or penetrating. Globally, trauma is a significant cause of morbidity and mortality [1]. Hollow viscus injuries mostly result from penetrating abdominal trauma. In blunt abdominal trauma, attention is usually more focused on the intra-abdominal solid organs and patient's hemodynamic status. Blunt abdominal trauma causing perforation of a hollow viscus is relatively an uncommon occurrence in children and diagnosis may not be prompt [2, 3]. Late presentation and treatment may result in peritonitis, increased morbidity and mortality. Prompt recognition and treatment is the cornerstone of management of hollow viscus injury [4]. Blunt abdominal trauma resulting in hollow viscus injury has been quoted to have an incidence of 1% to 8.5% and may vary with geographical location [5]. One study conducted in Brazil reported an incidence of 4 to 15% [6]. Hollow viscus injury necessitates operative treatment while most blunt abdominal trauma with solid organ injuries are managed non-operatively [7]. Limited experience on the management of hollow viscus injury results from uncommon nature of the condition. There is paucity of data on hollow viscus injury resulting from blunt abdominal trauma in Nigeria. Hence, there is need for a study to determine the incidence, etiology, management and outcome of children who sustained hollow viscus injury. The aim of this study was to evaluate our experience in management of children who sustained hollow viscus injury from blunt abdominal trauma.

2. Methods

This was a retrospective study of children, less than 15 years of age, who were managed for hollow viscus injury between January 2009 and December 2018 at the pediatric surgery unit of Enugu State University Teaching Hospital (ESUTH) in Enugu, Nigeria. ESUTH is a teaching hospital in Enugu, Nigeria. The hospital serves the people of Enugu State, which, according to the 2016 estimates of the National Population Commission and the Nigerian National Bureau of Statistics, has a population of about four million and a population density of 616.0/km². The hospital also receive referral from peripheral



hospitals. Patients who have had surgery for a hollow viscus injury at a peripheral hospital before referral to ESUTH for reoperation were excluded from the study. The information extracted from the patients’ medical records included age of the patient, gender, interval between the incident and presentation, mechanism (etiology) of injury, clinical state at presentation, presence/absence of associated intra-abdominal injury, presence/absence of peritonitis, associated extra-abdominal injuries, intra-operative finding, operative procedure performed, complications of treatment, duration of hospital stay and outcome of treatment. The period of follow up was for 12 months.

Gender	Percentage
Male	15 (71.4%)
Female	6 (28.6%)
Median age of the patients	10 years (8-14)
Mean interval between incident and presentation	5 days (3 – 10)
Presented within 24 hours	2 (9.5%)
Presented between 24 and 48 hours	3 (14.3%)
Presented after 48 hours	16 (76.2%)
The mean duration of hospital stay	16 days (11-25)

Table 1: Patients’ demographic features (N=21)

Ethical approval was obtained from the research ethics committee of ESUTH. Patients’ caregivers gave their consent for the study. Statistical Package for Social Science (SPSS) for Windows version 23 (IBM Corp., Armonk, NY) was used for data entry and analysis. Data were expressed as percentages, median, mean, and range.

3.2. Mechanism (etiology) of injury.

Road traffic accident was the most common mechanism of injury. Others are depicted in Table 2.

Pre-operative protocol

On presentation to the hospital, the patients were clinically evaluated and resuscitated using Advanced Trauma Life Support (ATLS) protocol. Abdominal ultrasound and radiographs were performed and further imaging investigation such as computed tomography (CT) scan was requested based on findings on clinical assessment. Urgent hemoglobin estimation and serum electrolytes were obtained. Anemia and electrolyte derangements (if any) were corrected and patient taken to theater for surgery. All the patients had laparotomy under general anesthesia.

Mechanism of injury	Number of patients	Percentage
Road traffic accident		
Pedestrian	5	23.8
Passenger	3	14.3
Fall from a height	7	33.3
Assault	3	14.3
Recreational accident	2	9.5
Child abuse	1	4.8

Table 2: Mechanism of injury (N=21)

Intra-operative protocol

Abdominal access was through transverse abdominal incision. An abdominal exploration was performed and any damaged hollow viscus was repaired as appropriate. The definitive surgery performed was dependent on the particular viscus that was damaged. The surgical procedures were carried out by a pediatric surgeon.

3.3. Symptoms at presentation

The patients presented with abdominal pain, vomiting, constipation and abdominal distension in various combinations: Abdominal pain 21 (100%); vomiting 13 (61.9%); constipation 11 (52.4%) and abdominal distension 5 (23.8%).

Post-operative protocol

Oral intake was started on return of bowel function. Post-operative complications were documented. Discharge was based on full establishment of oral intake and adequate wound healing.

3.4. Investigation performed

All the patients had abdominal/chest x ray and abdominal ultrasound. Eleven (52.4) patients showed free air under the diaphragm on radiograph. Abdominal ultrasound showed the presence of echo rich peritoneal fluid in 15 (71.4%) patients. Computed tomography (CT) scan was performed in 5 (23.8%) patients.

3. Results

3.1. Demographic profile of the patients

A total of 583 cases of blunt abdominal traumas were recorded during the study period, out of which 21 cases had hollow viscus injury. This translates to 3.6 percent. Details of the patients demography is shown in Table 1.

3.5. Hemodynamic status/peritonitis at presentation

Six (28.6%) patients were in a clinical state of shock at presentation. This was evidenced by tachycardia, hypotension and narrow pulse pressure. Five (23.8%) patients had peritonitis which was characterized by rebound tenderness, fever, guarding and rigidity.

3.6. Associated intra-abdominal/extra-abdominal injury

Three (14.3%) patients had associated splenic injury while two (9.5%) patients had associated liver injuries. One (4.8%) patient each had an associated head injury, right femoral fracture and left hemothorax.



3.7. Intra-operative finding and operative procedure performed

The small intestine was the most injured organ. Others are shown in Table 3.

Intra-operative finding	Operative procedure performed	Number (%)
Small intestinal perforation		
Ileum	Primary repair	10 (47.6)
Jejunum	Primary repair	6 (28.6)
Urinary bladder rupture	Primary repair	2 (9.5)
Mesenteric injury with bowel gangrene	Resection and anastomosis	2 (9.5)
Large bowel perforation	Repair + ileostomy	1 (4.8)

Table 3: Intra-operative finding and procedure performed (N=21)

3.8. Complications of treatment

There was no complication in 11 (52.4%) patients. Surgical site infection was the most common post-operative complication as reflected in Figure 1.




Figure 1: Post-operative complication (N=21)

3.9. Outcome of management

Nineteen (90.5%) patients recovered and were discharged home. One (4.8%) patient signed out against medical advice. One (4.8%) patient who had post-operative enterocutaneous fistula expired.

Early recognition of hollow viscus injury may not be easy and presents a task to the managing team [8]. The most frequent missed injury in blunt abdominal trauma is intestinal injury [9]. Forces such as compression or deceleration forces are responsible for hollow viscus injury. Compression force increases the intraluminal pressure resulting in rupture. Vascular injuries result from deceleration forces causing stretching/tearing of the mesentery [10]. Hollow viscus injury ranges from minor hematoma to frank perforation; it could also be partial thickness or full thickness injury.

In the present study, the incidence of hollow viscus perforation is consistent with the report of a similar study conducted in Turkey [11]. However, an incidence as low as 1% and as high as 15% has been reported [6, 12]. The difference in incidence may be due to the mechanism or severity of the trauma. The male predominance recorded in the current study is in line with the report of other studies [11, 13]. The exact reason for the male dominance is not clear but may be due to the fact that males behave more impulsively and take more risks than females. The median age of our patients is in agreement with the findings of other researchers [13, 14]. Howbeit, Borgianni et al reported a median age of 12 years [15]. The discrepancies in the median age of the patients in different studies may be explained by predominant etiology of the injury. For instance, in hollow viscus injuries resulting from abuse, the median age of the patients was 2.5 years [16]. The mean interval before presentation of our patients is not in agreement with the finding of Jha et al [17]. The late presentation of the patients may be due to ignorance and poverty which are quite common in low-income countries. There are wide variations in the duration of hospital stay of patients who sustain hollow viscus [13, 18, 19]. The duration of stay may be dependent on the extent of operative procedure and presence of associated injuries. Teuben et al reported longer stay in patients who have associated injuries [20]. Road traffic accident was the common mechanism of injury in the present study. This is comparable to the report of other series on hollow viscus injury [11, 19]. In developing countries, fall from height is a significant cause of hollow viscus injury [17, 21]. The explanation for the fall from height in children in low-income countries could be due to children climbing trees to pluck fruits. One uncommon cause of hollow viscus injury in children is abuse. Child abuse should be suspected in hollow viscus injury when the child or caregiver is in denial or when the degree of trauma is disproportionate to the extent of clinically observed injury [12]. The clinical presentation of children with hollow viscus injury may depend on the part of bowel that is injured and interval between the time of the incident and time of presentation to the hospital. Features of peritonitis may develop as early as 48 hours after a hollow viscus injury [13].

About half of our patients showed free air under the diaphragm. This is at variance with a report from South Asia [17]. The time interval between the incident and taking of the radiograph may explain the presence or absence of air under the diaphragm since it may take some time for air to accumulate in the peritoneal cavity. Abdominal ultrasound is essential for the detection of the site of hollow viscus perforation and echo rich peritoneal fluid. The ability of the sonographer to detect these features may depend on his experience and expertise. Computed tomography (CT) scan is



considered the gold standard for evaluation of hollow viscus injury, however, non-availability, cost and exposure to significant radiations are the problems associated with CT scan [22].

About 29% of our patients presented in shock. One study 4. conducted in Zaria, Nigeria reported that 23% of their patients presented in shock [23]. The late presentation of our patients may have accounted for the higher number of patients in shock at 5. presentation. The associated injuries in our patients are similar to the findings of a study from Jharkhand, India [17]. Mortality is increased when hollow viscus injury is associated head injury [13]. 6. The small bowel is the most injured hollow viscus in the index study. This finding is a consistent observation by other researcher [11, 17]. Gastric and urinary bladder injuries following blunt 7. abdominal trauma have also been reported [24, 25]. The exact factors that determine which organs are injured in blunt abdominal trauma are unknown although the intensity of the trauma and area of impact may play a role. Surgical site infection following 8. laparotomy for injured viscus is a common post-operative complication [26, 27]. This may be due to the bacterial loads in the intestine. Enterocutaneous fistula may follow anastomotic leak when anastomosis is performed in a background of gross 9. peritoneal contamination [17].

The mortality rate of 4.8% recorded in the present study is comparable to the report of Jha et al. [17] However, Gunturi et al reported a mortality rate of 15% [28]. Mortality following surgery for hollow viscus injury could be multifactorial: disease related factors, patient related factors or surgeon related factors.

5. Conclusion

Hollow viscus injury following blunt abdominal trauma is uncommon in children and the small intestine was the most injured. In abdominal trauma following blunt abdominal trauma, a high index of suspicion for hollow viscus injury is required to avoid late diagnosis and its attendant morbidity/mortality. Emphasis should be placed on prevention which can be achieved by good parental control.

Limitation of the study

1. Children who succumbed to their hollow viscus injury before presentation to the hospital were not captured in this study.
2. Small sample size. A larger number of patients would have availed better analysis.

Declaration of conflict of interest: The author declares that there is no conflict of interest.

Sources of support: None

References

1. Sakran JV, Greer SE, Werlin E et al (2012) Care of the injured worldwide: trauma still the neglected disease of modern society. *Scand J Trauma Resusc Emerg Med.*; 20: 64.
2. Ozturk H, Onen A, Otcu S et al (2003) Diagnostic Delay Increases Morbidity in Children with Gastrointestinal Perforation from Blunt Abdominal Trauma.; 33: 178-182.
3. Fakhry SM, Brownstein M, Watts DD et al (2000) Relatively short diagnostic delays (<8 hours) produce morbidity and mortality in blunt small bowel injury: an analysis of time to operative intervention in 198 patients from a multicenter experience. *J Trauma.*; 48(3): 408-415.
4. Matsushima K, Mangel PS, Schaefer EW et al. Blunt hollow viscus and mesenteric injury: still underrecognized. *World J Surg.* 2013; 37(4): 759-765.
5. Canty TG Sr, Canty TG Jr, Brown C. Injuries of the gastrointestinal tract from blunt trauma in children: a 12-year experience at a designated pediatric center. *J Trauma.* 1999; 46(2): 234-240.
6. Brusca V, Coimbra R, Rasslan S et al. Blunt gastric injury. A multicentre experience. *Injury.* 2001; 32(10): 761-764.
7. Swaid F, Peleg K, Alfici R et al. Concomitant hollow viscus injuries in patients with blunt hepatic and splenic injuries. An analysis of a National Trauma Registry database. *Injury.* 2014; 45(9): 1409-1412.
8. Gonser-Hafertepen LN, Davis JW, Bilello JF et al. Isolated free fluid on abdominal computed tomography in blunt trauma: watch and wait or operate? *J Am Coll Surg.* 2014; 219(4): 599-605.
9. Lawson CM, Daley BJ, Ormsby CB et al. Missed injuries in the era of the trauma scan. *J Trauma.* 2011; 70(2): 452-458.
10. Kulvatunyou N, Albrecht RM, Bender JS et al. Severe abdominal wall disruption, hollow viscus injury and major vascular injury. *Am Surg.* 2011; 77(5): 534-538.
11. Ciftci AO, Tanyel FC, Salman AB et al. Gastrointestinal tract perforation due to blunt abdominal trauma. *Pediatr Surg Int.* 1998; 13(4): 259-264.
12. Aguirre J, Perez L, Retamal A et al. Gastrointestinal injuries from blunt abdominal trauma in children. *Rev Chil Radiol.* 2014; 20(3): 105-111
13. Abbas SM, Upadhyay V. Hollow viscus injury in children: Starship Hospital experience. *World J Emerg Surg.* 2007; 2: 14.
14. Holmes JF, Lillis K, Monroe D et al. Identifying children at a very low risk of clinically important blunt abdominal injuries. *Ann Emerg Med.* 2013; 62(2): 107-116.e2.
15. Borgialli DA, Ellison AM, Ehrlich P et al. Association between seat belt sign and intra-abdominal injuries in children with blunt torso trauma in motor vehicle collisions. *Acad Emerg Med.* 2014; 21(11): 1240-1248.
16. Muniz A. Evaluation and Management of Pediatric Abdominal Trauma. *Ped Emerg Med Pract.* 2008; 5(3): 1-32
17. Jha NK, Yadav SK, Sharma R et al. Characteristics of Hollow Viscus Injury following Blunt Abdominal Trauma: a Single Centre Experience from Eastern India. *Bull Emerg Trauma.* 2014; 2(4): 156-160
18. Lane WG, Lotwin I, Dubowitz H et al. Outcome for children hospitalized with abusive versus noninflicted abdominal trauma. *Pediatrics.* 2011; 127(6): e1400-e1405.
19. Mingoli A, La Torre M, Brachini G et al. Hollow viscus injury: predictors of outcome and role of diagnostic delay. *Therapeutics and clinical risk management.* 2017; 13: 1069-1076.
20. Teuben MPJ, Spijkerman R, Blokhuis TJ. Safety of selective nonoperative management for blunt splenic trauma: the impact of concomitant injuries. *Patient Saf Surg.* 2018; 12: 32.
21. Chirdan LB, Uba AF, Chirdan OO.. Gastrointestinal injuries following blunt abdominal trauma in children. *Niger J Clin*



- Pract. 2008; 11(3): 250-253.
22. Cinquantini F, Tugnoli G, Piccinini et al. Educational Review of Predictive Value and Findings of Computed Tomography Scan in Diagnosing Bowel and Mesenteric Injuries After Blunt Trauma; Correlation With Trauma Surgery Findings in 163 Patients. *Can Assoc Radiol J.* 2017; 68(3): 276-285.
 23. Ameh EA, Nmadu PT. Gastrointestinal injuries from blunt abdominal trauma in children. *East Afr Med J.* 2004; 81(4): 194-197.
 24. Aboobakar MR, Singh JP, Maharaj K et al. Gastric perforation following blunt abdominal trauma. *Trauma Case Rep.* 2017; 10: 12-15.
 25. Elkbuli A, Ehrhardt JD, Hai S. et al. Management of blunt intraperitoneal bladder rupture: Case report and literature review. *Int J Surg Case Rep.* 2019; 55: 160-163.
 26. Jain S, Maske D, Songra MC. Clinical study of hollow viscus injury in abdominal trauma. *Int Surg J.* 2018; 5(1): 39-44.
 27. Gandhi A, Sharma R, Sinha DK. Characteristics of hollow viscus injury following blunt abdominal trauma: A study done in Rajendra institute of medical sciences. *International Journal of Surgery Science.* 2019; 3(2): 135-138.
 28. Gunturi SRV, Thumma VM, Bathalapalli JMR et al. Hollow viscus injury due to blunt abdominal trauma. *International Surgery Journal.* 2017; 4(3): 861-865.