

Studies the Residual Effect of Six Herbicides Applied to The Minimum Tillage Non-Puddled Transplanted Winter and Summer Rice in Bangladesh

Mohammad Mobarak Hossain *, Mahfuza Begum and Md. Moshir Rahman
Department of Agronomy, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh

Article Info

Received: February 21, 2022

Accepted: March 08, 2022

Published: April 22, 2022

***Corresponding author:** Mohammad Mobarak Hossain, Department of Agronomy, Bangladesh Agricultural University, Mymensingh 2202, Bangladesh.

Citation: Mohammad Mobarak Hossain, Mahfuza Begum and Md. Moshir Rahman. (2022) "Studies the Residual Effect of Six Herbicides Applied to The Minimum Tillage Non-Puddled Transplanted Winter and Summer Rice in Bangladesh.", Journal of Agricultural Research Pesticides and Biofertilizers, 3(4); DOI:<http://doi.org/03.2022/1.1068>.

Copyright: © 2022 Mohammad Mobarak Hossain. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

The poisonous effect of herbicide residues on the succeeding crops is one of the principal concerns against the safe use of herbicides for controlling weeds. Acute labor crisis for crop production pushes farmers to adopt minimum tillage non-puddled (MTNP) rice cultivation in Bangladesh. This on-farm research on plant bioassay was conducted at the Mymensingh region of Bangladesh during October–December in 2016 and 2017 year. Here, we studied any residual effect of six herbicides *viz.*, glyphosate, pendimethalin, ethoxysulfuron-ethyl, isoproturon, fenoxaprop-p-ethyl, and carfentrazone-ethyl + isoproturon on the succeeding mustard in a winter rice-summer rice-mustard rotation. These herbicides were applied in 10 different combinations for controlling weeds of MTNP winter and summer rice for two years. Immediately after the harvest of summer rice, the indicator crop mustard was planted on the same plots of the respective treatments of the previous MTNP rice experiment. Data of a two-year experiment revealed that, after 25 days of planting, the plant population, length of seedling root and shoot, and seedling biomass did not vary significantly across the treatments. Moreover, leaf chlorophyll content in all the treatments was statistically identical. Furthermore, an excellent growth mustard plant without any sign of phytotoxicity was observed in all treatments. Hence, it could be concluded that herbicides used for controlling weeds in MTNP rice did not retain any residues in soil enough to hamper the growth and development of succeeding mustard in winter rice-summer rice-mustard rotation Bangladesh.

Keywords: bioassay; herbicide residue; toxicity; chlorophyll; phytotoxicity

Introduction:

In Bangladesh, rice (*Oryza sativa* L.) is conventionally cultivated by transplanting seedlings into puddled soil, typically for ease of crop establishment and weed control [1]. But rice can be grown by transplanting them into minimum tillage non-puddled (MTNP) soils without any yield penalty but with additional benefits of saving land preparation costs, fuel energy requirement, labor, and irrigation water [2, 3]. However, severe weed infestation has been argued against the widespread adoption of MTNP rice cultivation [4, 5]. As a result, farmers are advised to do hand weeding up to six times throughout the rice-growing season to maintain weeds below the economic threshold level in this practice [6]. Due to the agricultural laborers' crisis and high wage rate, herbicides are being quickly embraced in countries with a manpower shortage for weeding [7, 8]. Recent advancements in broad-spectrum herbicides may offer an opportunity to manage weeds more successfully in non-puddled rice transplanting systems [9].

Unfortunately, the repeated use of these chemicals may lead to persists residues of herbicides in the soil [10]. Wyk and Reinhardt [11] discovered an excessive quantity of imazethapyr residue harmed corn planted after soybean. If herbicide residues remain in the soil, they may decrease the performance of subsequent crops [12]. Sulfonylurea herbicide residues even at low concentrations in soil may damage rotating crops [13, 14]. However, farmers often apply herbicides without understanding or evaluating the herbicide's residual impact on following crops.



Furthermore, minimal study on the residual effects of herbicides located at the Mymensingh region of Bangladesh (N: 24°75', E: 90°50') from October–March in 2016 and 2017 years under the general, a soil chemical test or bioassay may be performed to assess the residual herbicide content in the soil [15]. However, chemical analysis is prohibitively expensive, and therefore a plant bioassay in the field may be used to predict the presence of herbicides in soil. While a bioassay does not quantify the herbicide residue in the soil but shows whether there is enough residue in the soil to harm a succeeding crop practically. The bioassay in a similar field of previous herbicide-treated crops is more convenient and indicates the residual impact of herbicides in the field scenario.

Materials and Methods:

Location and tenure:

This on-farm experiment was conducted at the farmers' field at the farmers' field immediately after harvest of MTNP summer rice to determine the residual impact of rice herbicides on subsequent mustard under winter rice- summer rice-mustard cropping system in Bangladesh.

Soil condition:

The field was a well-drained medium medium-high land with sandy clay loam soil with sand, silt, and Clay @ 50, 23, and 27%, respectively, and soil pH of 7.2.

Climatic statement:

The region gets an average annual rainfall of 178 millimeters, with about 93% of it falling between May and September (Figure 1). Total rain was greatest during the summer rice season and lowest during the winter rice season in both years. Occasionally, the highest average temperature was about 29°C in April–May, while the lowest temperature was approximately 13°C in January. In both years, the months of October–November, and March had the most sunshine hours.




Figure 1: Climatic condition of the on-farm experimental site at the Mymensingh region of Bangladesh during 2016 and 2017.

MTNP rice experiment:

Winter rice (*Oryza sativa* L.) during January–May and summer rice during June–September was grown under MTNP system continuously two years during 2016 and 2017. The MTNP rice herbicides were applied in winter and summer rice per the experiment used a four-replicated randomized complete block recommended rates and time (Table 2).

design. The unit plots were 9 m × 5 m in size. The MTNP land was prepared in a single pass operation, using the Versatile Multi-crop Planter (VMP) machine. Six rows each of 6 cm broad and 5 cm deep was made at a time. Total ten combinations (Table 1) of six continuously two years during 2016 and 2017.

Table 1: Treatments using different herbicides in MTNP rice experiments for weed control

Legends	Treatments
T ₀	Control
T ₁	Glyphosate
T ₂	Glyphosate <i>fb</i> Pendimethalin
T ₃	Glyphosate <i>fb</i> Ethoxysulfuron-ethyl
T ₄	Glyphosate <i>fb</i> Carfentrazone-ethyl+Isoproturon
T ₅	Glyphosate <i>fb</i> Isoproturon



T ₆	Glyphosate <i>fb</i> Fenoxaprop-p-ethyl
T ₇	Glyphosate <i>fb</i> Pendimethalin <i>fb</i> Ethoxysulfuron-ethyl
T ₈	Glyphosate <i>fb</i> Pendimethalin <i>fb</i> Carfentrazone-ethyl+Isoproturon
T ₉	Glyphosate <i>fb</i> Pendimethalin <i>fb</i> Isoproturon
T ₁₀	Glyphosate <i>fb</i> Pendimethalin <i>fb</i> Fenoxaprop-p-ethyl

Table 2: Statements of herbicides used in MTNP rice

Name of herbicides	Chemical group	Rate (a.i. ha ⁻¹)	Applied at	Field condition
Glyphosate	Phosphonic acid	9 L	3 DBP	Field capacity
Pendimethalin	Dinitroaniline	11 L	3 DAP	Field capacity
Ethoxysulfuron-ethyl	Sulfonylurea	667 g	25 DAP	Saturation
Carfentrazone-ethyl	Triazolinone	2.5 kg	25 DAP	Field capacity
Isoproturon	Urea	3.4 L	25 DAP	Field capacity
Fenoxaprop-p-ethyl	Aryloxyphenoxy-propionate	7.2 L	25 DAP	Field capacity

*a.i. = active ingredient, DBP = Days before planting, DAP = Days after planting

Bioassay experiment:

Measurements:

The research used the bioassay technique to determine the residual impact of herbicides applied to winter and summer rice on mustard (*Brassica napus* L.). Mustard was grown from October–December. On the relevant plots, planting of 7 kg, ha⁻¹ three quadrates was reported. The length of root and shoot and the seeds were done using VMP on October 05 in 2016 and 2017, immediately after the harvest of summer rice. Light watering was done after planting for optimal germination. During the residual impact research period, no fertilizer was applied in the field. Before planting in experimental plots, sample seed germination was examined in the laboratory, and >80% germination was reported. Weeds were maintained below the economic threshold level by manual weeding throughout the growing season [16].

A 1.0 m × 1.0 m quadrat was used to record the plant population on m². The quadrat was randomly put in three locations within each plot. Plants were counted inside the quadrat, and an average of three quadrates was reported. The length of root and shoot and the biomass of 25 days aged seedlings was determined by examining the biomass after 72 hours of drying at 70°C of randomly selected ten plants. The chlorophyll content of the leaves using SPAD 502 Plus Chlorophyll Meter from the young, tender leaf of these plants. Phyto-toxicity of herbicide on mustard and crop vigor of mustard was assessed visually four times at 15 days of interval up to 60 DAS following the toxicity grading of IRRI [17] (Table 3) and crop vigor scale [18] as of; 1: Poor, 2: Fair, 3: Good and 4: Excellent.

Table 3: Phyto-toxicity scoring

Toxicity	Rating
Normal growth: non-toxic	1
Slightly toxic: Injury/discoloration recoverable	2
Moderately toxic: Some stunting/discoloration recoverable	3
Severely toxic: Stand loss irrecoverable	4
Toxic (Plant kill): Total damage	5

Data analysis:

The data were analyzed using the statistical software *STAR* following analysis of variance, and treatment means were separated using the Duncans' Multiple Range Test at a significance level of 5%. The plant population of mustard m⁻² areas at 25 DAS did not vary significantly ($p>0.05$) at both 2016 and 2017 years by the residues of six MTNP rice herbicides (Figure 2). We recorded >80 plants m⁻² across all the treatments in both years. Data implies that the previously applied herbicides did persist in the soil enough to hamper the germination capacity of mustard.

Results:

Effect on plant population:




Figure 2: Residual effect of herbicides on the plant population of mustard at $p > 5\%$ level. T₀ = Control, T₁ = Glyphosate, T₂ = Glyphosate followed by (fb) Pendimethalin, T₃ = Glyphosate fb Ethoxysulfuron-ethyl, T₄ = Glyphosate fb Carfentrazone-ethyl+Isoproturon, T₅ = Glyphosate fb Isoproturon, T₆ = Glyphosate fb Fenoxaprop-p-ethyl, T₇ = Glyphosate fb Pendimethalin fb Ethoxysulfuron-ethyl, T₈ = Glyphosate fb Pendimethalin fb Carfentrazone-ethyl+Isoproturon, T₉ = Glyphosate fb Pendimethalin fb Isoproturon, T₁₀ = Glyphosate fb Pendimethalin fb Fenoxaprop-p-ethyl

Effect on the length of root and shoot at 25 DAS:

Data presented in Table 4 indicated a statistically non-significant ($p > 5\%$) effect of previously used herbicides on the root and shoot development of succeeding lengths of mustard at 25 DAS in both 2016 and 2017 year. None

of the other treatments relative to Control did inhibit the root and shoot growth. Results revealed no residual effect of MTNP rice

Table 4: Residual effect of herbicides on the root and shoot length of mustard

Treatments	Root length (cm)		Shoot length (cm)	
	2016	2017	2016	2017
Control	6.41	5.26	16.82	18.5
Glyphosate	6.45	5.32	22.01	22.5
Glyphosate fb Pendimethalin	7.39	5.68	22.72	20.6
Glyphosate fb Ethoxysulfuron-ethyl	7.42	5.58	21.30	24.0
Glyphosate fb Carfentrazone-ethyl+Isoproturon	6.62	6.06	21.22	20.5
Glyphosate fb Isoproturon	7.42	5.91	20.55	22.5
Glyphosate fb Fenoxaprop-p-ethyl	6.39	5.88	22.71	23.5
Glyphosate fb Pendimethalin fb Ethoxysulfuron-ethyl	6.45	6.32	22.23	25.0
Glyphosate fb Pendimethalin fb Carfentrazone-ethyl+Isoproturon	7.43	5.93	18.61	21.0
Glyphosate fb Pendimethalin fb Isoproturon	6.56	5.97	20.27	23.3
Glyphosate fb Pendimethalin fb Fenoxaprop-p-ethyl	7.47	5.87	20.93	20.5
Standard deviation	0.09	0.30	2.06	1.83
Co-efficient of variance	1.25	5.27	10.32	8.36
Standard error of mean difference	0.03	0.09	2.98	2.41

Effect on the leaf chlorophyll content at 25 DAS:

The chlorophyll contents of mustard leaves based on the SPAD meter reading reported a non-significant ($p > 5\%$) variation by the

carryover effect of rice herbicides (Figure 3). Data reveal that six herbicides in 10 combinations used in earlier MTNP winter and

summer rice to control weeds does not persist in the soil to hamper the leaf chlorophyll contents of succeeding mustard.




Figure 3: Residual effect of herbicides on the chlorophyll content of mustard leaves at $p > 5\%$ level. T₀ = Control, T₁ = Glyphosate, T₂ = Glyphosate followed by (fb) Pendimethalin, T₃ = Glyphosate fb Ethoxysulfuron-ethyl, T₄ = Glyphosate fb Carfentrazone-ethyl+Isoproturon, T₅ = Glyphosate fb Isoproturon, T₆ = Glyphosate fb Fenoxaprop-p-ethyl, T₇ = Glyphosate fb Pendimethalin fb Ethoxysulfuron-ethyl, T₈ = Glyphosate fb Pendimethalin fb Carfentrazone-ethyl+Isoproturon, T₉ = Glyphosate fb Pendimethalin fb Isoproturon, T₁₀ = Glyphosate fb Pendimethalin fb Fenoxaprop-p-ethyl

Effect on seedling biomass of mustard at 25 DAS:

Statically similar ($p > 5\%$) biomass of the 25 days aged seedlings of mustard at all the treatments relative to Control was recorded in both years in this study (Table 5).

shown), indicating an excellent crop growth without any poisonous symptoms relative to Control treatment. Such results suggest no carryover effect of previously used six MTNP rice herbicides on the succeeding mustard.

Discussion:

The current two-year on-farm research investigated any potential persistence impact of six rice herbicides (glyphosate, pendimethalin, ethoxysulfuron-ethyl, carfentrazone-ethyl + isoproturon, isoproturon, and fenoxaprop-p-ethyl) on the indicator crop plant mustard. The results demonstrated applied rice herbicides in 10 different combinations had no harmful effect on subsequent mustard plant population, length of shoot and root, leaf chlorophyll content and seedling biomass. Moreover, no toxic symptoms were observed visually on healthy plant growth across all the treatments. The finding of prior research agrees with our results showing that herbicides used in the preceding wheat crop did not affect maize germination [19, 20]. Additionally, they noticed no apparent phototoxicity on mustard by the residues of imazethapyr + pendimethalin applied to black gram. Another research has shown that herbicides applied to onions [21] and peanuts [22] did not substantially impact the germination of subsequent sorghum and wheat, and gram. Khokhar and Charak [23] also found that herbicides sprayed to wheat had no discernible impact on the germination of the subsequent maize, green gram, and cucumber. The explanation for this may be linked to the degradation of all herbicides in soil [24] which is related to the half-life of the herbicides examined. For example, half-life (days) of glyphosate: 30–32 days [25], pendimethalin: 25–35 days [26], ethoxysulfuron: 60 days [27], carfentrazone-ethyl: 3.8–5.8 hours only [28], isoproturon: 24 days [29] and fenoxaprop-p-ethyl: 1.45–2.30 days [30]. On the other hand, mustard required about 90 days to harvest in our prior study. Thus, as Parthipan et al. [31] and Yazdanpak et al. [32] indicated, there was little chance of these herbicides persisting in the soil until the next crop growing season.

Table 5: Residual effect of herbicides on seedling biomass of mustard at 25 DAS

Treatments	Plant biomass (g 10 plant ⁻¹)	
	2016	2017
Control	1.40	1.31
Glyphosate	1.47	1.32
Glyphosate fb Pendimethalin	1.46	1.29
Glyphosate fb Ethoxysulfuron-ethyl	1.43	1.36
Glyphosate fb Carfentrazone-ethyl+Isoproturon	1.41	1.42
Glyphosate fb Isoproturon	1.46	1.40
Glyphosate fb Fenoxaprop-p-ethyl	1.47	1.39
Glyphosate fb Pendimethalin fb Ethoxysulfuron-ethyl	1.46	1.43
Glyphosate fb Pendimethalin fb Carfentrazone-ethyl+Isoproturon	1.51	1.47
Glyphosate fb Pendimethalin fb Isoproturon	1.53	1.49
Glyphosate fb Pendimethalin fb Fenoxaprop-p-ethyl	1.42	1.41
Standard deviation	0.12	0.11
Co-efficient of variance	8.24	7.34
Standard error of mean difference	2.27	2.11

Visual scoring of phytotoxicity and crop vigor:

The visual observation scoring of toxicity symptoms on the morphology scored "1" and crop vigor scored "4" (data not



The unaffected germination rate might have influenced the to obtain a similar plant population of succeeding mustard across all treatments in this study.

At 25 DAS, the current research discovered no significant impact on the seedlings' shoot and root length and dry matter production. This finding supports Taslima et al. [33]. They disclosed no adverse effects of the residues of eight herbicides (pendimethalin, pretilachlor, triasulfuron, ethoxysulfuron, pyrazosulfuron-ethyl, carfentrazone-ethyl, 2,4-D amine, and carfentrazone-ethyl + isoproturon) on the biomass of succeeding mungbean, sunflower, and jute. The research findings of Yadav and Bhullar [34] also discovered that herbicides applied to soybean had no impact on the dry matter buildup of succeeding wheat, barley, spinach, pea, raya, canola, and sugarbeet due to thoroughly degradation of prior applies herbicides. Further research by Sangeetha et al. [35], Bahrapor and Ziveh [36], and Yadav et al. [37] confirmed no significant residual toxicity in shoot length was seen in the following soybean and wheat treated with herbicides in the prior crop. Similarly, Rathod et al. [21] found that onion herbicide residue had no detrimental effect on the dry matter accumulation of the following sorghum.

Herbicide applied to the MTNP rice did not affect the chlorophyll content of the indicator crop mustard leaves in this research. The excellent plant growth resulted in increased leaf area facilitated to have a higher efficiency of light, water, and nutrients use [38], resulting in increased plant biomass both in the herbicidal and Control treatment in this study. Prior studies assert that herbicide residue had no detrimental impact on phenotypic and genotypic development resulting from the following crops' regular leaf chlorophyll content [31, 35]. Taslima et al. [33] also found an unaffected chlorophyll content in leaves of succeeding sunflower, mungbean, and jute when eight different herbicides were applied to prior wheat.

Any non-toxic effect of herbicide residue on the length of root, leaf chlorophyll content, and plant dry matter might have influenced the non-persistent herbicides in soil. Applied herbicides in may be broken down by the cultural activities of various crops, such as flooding for irrigation and microbial degradation, are the primary mechanisms by which herbicides are dissipated from the soil [39, 40]. Thus, one might argue that many herbicides used for weed management are safe in terms of residual toxicity in soil [41, 42]. The explanation for this may be because the herbicides used have entirely degraded in the soil or that their presence is at a measurable level that does not negatively impact the growth of subsequent crops. Previous investigations concluded that residues of the majority of herbicides remained below the detectable level in the soil after 30–120 days of treatment [26, 43]. Hence, the above-discussed reasons clarify those six herbicides in ten combinations tested in MTNP rice pose no detrimental residual impact on the growth and development of subsequent mustard in Bangladesh.

Conclusion:

The results indicated that ten combinations of six herbicides: glyphosate, pendimethalin, ethoxysulfuron-ethyl, isoproturon, fenoxaprop-p-ethyl, and carfentrazone-ethyl + isoproturon applied to MTNP winter and summer rice had no toxic effect on the plant population, seedling growth in terms of root and shoot length, and

biomass and leaf chlorophyll content of succeeding mustard with excellent growth without any phytotoxic symptoms. Thus, the study concluded that herbicides used in preceding MTNP rice are safer for the next seasons' crop cultivation in rotation.

Reference:

- Bell, R. W., Haque, M., Jahiruddin, M., Rahman, M., Begum, M., Miah, M. A., Islam, M., Hossen, M., Salahin, N., Zahan, T., Hossain, M. M., Alam, M. K., & Mahmud, M. N. H. (2019). Conservation agriculture for rice-based intensive cropping by smallholders in the eastern gangetic plain. *Agriculture (Switzerland)*, 9(1).
- Gathala, M. K., Timsina, J., Islam, M. S., Rahman, M. M., Hossain, M. I., Harun-Ar-Rashid, M., Ghosh, A. K., Krupnik, T. J., Tiwari, T. P., & McDonald, A. (2015). Conservation agriculture based tillage and crop establishment options can maintain farmers' yields and increase profits in South Asia's rice-maize systems: Evidence from Bangladesh. *Field Crops Research*, 172, 85–98. <https://doi.org/10.1016/j.fcr.2014.12.003>
- Townsend, T. J., Ramsden, S. J., & Wilson, P. (2016). Analysing reduced tillage practices within a bio-economic modelling framework. *Agricultural Systems*, 146, 91–102.
- Sims, B., Corsi, S., Gbehounou, G., Kienzle, J., Taguchi, M., & Friedrich, T. (2018). Sustainable weed management for conservation agriculture: options for smallholder farmers. *Agriculture*, 8(8), 118.
- Nichols, V., Verhulst, N., Cox, R., & Govaerts, B. (2015). Weed dynamics and conservation agriculture principles: A review. *Field Crops Research*, 183, 56–68.
- Farooq, M., Flower, K. C., Jabran, K., Wahid, A., & Siddique, K. H. M. (2011). Crop yield and weed management in rainfed conservation agriculture. *Soil and Tillage Research*, 117, 172–183.
- Rashid, M. H., Alam, M. M., Rao, A. N., & Ladha, J. K. (2012). Comparative efficacy of pretilachlor and hand weeding in managing weeds and improving the productivity and net income of wet-seeded rice in Bangladesh. *Field Crops Research*, 128, 17–26.
- Dahal, S., & Bahadur Karki, T. (2014). Conservation agriculture based practices affect the weed dynamics in spring maize. *World Journal of Agricultural Research*, 2(6A), 25–33.
- Huang, J., Wang, S., & Xiao, Z. (2017). Rising Herbicide Use and its driving forces in China. *The European Journal of Development Research*, 29(3), 614–627.
- Tworokski, T. J., Welker, W. V., & Vass, G. D. (2000). Soil residues following repeat applications of Diuron, Simazine, and Terbacil. *Weed Technology*, 14(1), 191–196.
- Wyk, L. J. V., & Reinhardt, C. F. (2001). A Bioassay technique detects imazethapyr leaching and liming-dependent activity. *Weed Technology*, 15(1), 1–6.
- Houge, E., & Neilsen, G. (n.d.). Effects of excessive annual applications of ternacil, diuron, simazine and dichlobenil on vigor, yield and cation nutrition of young apple trees. *Candian Journal of Plant Science*, 68(3), 843–850.
- Janaki, P., Sharma, N., Chinnusamy, C., Sakthivel, N., & Nithya, C. (2015). Herbicide residues and their management strategies. *Indian Journal of Weed Science*, 47(3), 329–344.
- Hernández-Sevillano, E., Villarroya, M., Alonso-Prados, J. L.,



- & García-Baudín, J. M. (2001). Bioassay to detect MON-37500 and Triasulfuron residues in soils. *Weed Technology*, 15(3), 447–452.
15. Peachey, E. (2015). Testing for and Deactivating Herbicide Residues. In *Pacific Northwest Pest Management Handbooks* (2021st ed., pp. 25–32). OSU Extension Service - Extension and Experiment Station Communications.
 16. Hanson, B. D., & Thill, D. C. (2001). Effects of imazethapyr and pendimethalin on lentil (*Lens culinaris*), pea (*Pisum sativum*), and a subsequent winter wheat (*Triticum aestivum*) Crop. *Weed Technology*, 15(1), 190–194.
 17. IRRI. (1965). *Annual report for 1963*. Los Baños, Laguna, The Philippines.
 18. Kingman, A. R., & Moore, J. (1982). Isolation, purification and quantitation of several growth regulating substances in *Ascomyllum nodosum* (Phaeophyta). *Botanica Marina*, 25(4), 149–154.
 19. Kaur, T., & Brar Singh, S. (2014). Residual effect of sulfonylurea herbicides applied to wheat on succeeding maize-Indian Journals. *Indian Journal of Weed Science*, 46(2), 129–131.
 20. Kumar, S., Bhatto, M., Punia, S., & Rajni, P. (2015). Bioefficacy of herbicides in blackgram and their residual effect on succeeding mustard-Indian Journals. *Indian Journal of Weed Science*, 47(2), 211–213.
 21. Rathod, A., Solanki, R., Modhavadia, J., & Padamani, D. (2014). Efficacy of pre-and post-emergence herbicides in onion and their carry over effect on the succeeding crops. *Annals of Agricultural Research New Series*, 35(2), 209–216.
 22. Vaghasia, P. M., & Nadiyadhara, M. V. (2013). Effect of post-emergence herbicides in groundnut and its residual effect on succeeding crops. *International Journal of Forestry and Crop Improvement*, 4(2), 54–58.
 23. Khokhar, A., & Charak, A. (2011). Bio-efficacy of herbicides against complex weed flora in wheat and their residual effects on succeeding crops. *Journal Crop and Weed*, 7(2), 164–167.
 24. Mohamed, F., El-Nady, & Elsayed, B. B. (2013). Effect of phytotoxicity of pendimethalin residues and its bioremediation on growth and anatomical characteristics of *Cucumis sativus* and *Echinochloa crus-galli* plants. *Asian Journal of Crop Science*, 5, 222–237.
 25. Giesy, J., Dobson, S., & Solomon, K. (2000). Ecotoxicological risk assessment for roundup herbicide. *Reviews of Environmental Contamination and Toxicology*, 167, 35–120.
 26. Sireesha, A., Rao, P. C., Rao, P. V., Swapna, G., & Ramalakshmi, C. (2012). Residues of pendimethalin and oxyfluorfen in radish and their persistence in soil. *Journal Crop and Weed*, 8(2), 120–125.
 27. Sondhia, S., & Dixit, A. (2012). Bioefficacy and persistence of ethoxysulfuron in rice. *ORYZA- An International Journal on Rice*, 49(3), 178–182.
 28. Han, L., Xu, Y., Dong, M., & Qian, C. (2007). Dissipation and residues of carfentrazone-ethyl in wheat and soil. *Bulletin of Environmental Contamination and Toxicology*, 79(4), 445–447.
 29. Accinelli, C., Screpanti, C., & Vicari, A. (2005). Influence of flooding on the degradation of linuron, isoproturon and metolachlor in soil. *Agronomy for Sustainable Development*, 25(3), 401–406.
 30. Tandon, S. (2019). Degradation of fenoxaprop-p-ethyl and its metabolite in soil and wheat crops. *Journal of Food Protection*, 82(11), 1959–1964.
 31. Parthipan, T., Ravi, V., Subramanian, E., & Ramesh, T. (2013). Integrated weed management on growth and yield of transplanted rice and its residual effect on succeeding black gram. *Journal of Agronomy*, 12(2), 99–103.
 32. Yazdanpak, A., Amiri, A., Faghihi, K., & Karimian, N. A. (2014). The residual effect of herbicides on the germination and early growth of shiraz wheat cultivar in the development of healthy agricultural crops. *American-Eurasian Journal of Agricultural & Environmental Sciences*, 14(2), 161–164.
 33. Taslima, Z., Muktadir, M. A., Rahman, M. M., & Ahmed, M. M. (2018). Response of the succeeding crops as affected by the residue of herbicides applied in wheat in Old Brahmaputra Floodplain, Bangladesh. *Annals of Agrarian Science*, 16(4), 451–457.
 34. Yadav, R., & Bhullar, M. S. (2014). Residual effects of soybean herbicides on the succeeding winter crops. *Indian Journal of Weed Science*, 46(3), 305–307.
 35. Sangeetha, C., Chinnusamy, C., & Prabhakaran, N. (2012). Efficacy of imazethapyr on productivity of soybean and its residual effect on succeeding crops. *Indian Journal of Weed Science*, 44(2), 135–138.
 36. Bahrapor, T., & Ziveh, P. S. (2013). Effects of residue sulfonylurea herbicides on wheat. *International Journal of Agronomy and Plant Production*, 4(10), 2707–2713.
 37. Yadav, A., Malik, R., Punia, S. S., Mehta, R., Bir, D., Amarjeet, & Beuinder. (2004). Studies on carry-over effects of herbicides applied in wheat on the succeeding crops in rotation. *Indian Journal of Weed Science*, 36(1 & 2), 15–18.
 38. Divsalar, M., Oskouei, B., & Sheidaei, S. (2016). Study on correlation of seed germination percent of two sweet corn hybrids with field emergence and some measured traits related to yield. *Journal of Applied Environmental and Biological Sciences*, 6(3), 44–50.
 39. Lim, C. A. A., Awan, T. H., Cruz, P. C. S., & Chauhan, B. S. (2015). Influence of environmental factors, cultural practices, and herbicide application on seed germination and emergence ecology of *Ischaemum rugosum* Salisb. *PLOS ONE*, 10(9), e0137256.
 40. Jape, P., Maheshwari, V., & Chaudhari, A. (2019). Microbial Degradation of Nitroaromatic Pesticide: Pendimethalin. In D. P. Singh, V. K. Gupta, & R. Prabha (Eds.), *Microbial Interventions in Agriculture and Environment: Volume 1: Research Trends, Priorities and Prospects* (pp. 531–544). Springer.
 41. Aktar, Md. W., Sengupta, D., & Chowdhury, A. (2009). Impact of pesticides use in agriculture: Their benefits and hazards. *Interdisciplinary Toxicology*, 2(1), 1–12.
 42. Sangeetha, C., Chinnusamy, C., & Prabhakaran, N. (2011). Performance of early post-emergence herbicide in irrigated soybean (Glycine max (L.) Merrill). *Madras Agricultural Journal*, 98(4–6), 144–146.
 43. Zahan, T., Bell, R. W., Rahman, M., & Ahmed, M. M. (2020). Performance of pyrazosulfuron-ethyl in non-puddled transplanted rainy season rice and its residual effect on growth of the succeeding crop in rice-wheat cropping pattern. *International Journal of Pest Management*, 66(2), 122–130.