

Case Report: Cold Thermal Injury and Syncope from Pressurized Dust Cleaning Spray

James Espinosa MD^{1*}, Alan Lucerna DO², Risha Hertz APN³

¹Department of Emergency Medicine, Rowan University SOM/Jefferson - Stratford, NJ

²Program Director, Emergency Medicine, Jefferson NJ/Rowan University SOM, Stratford, NJ

³Independent researcher, Voorhees, NJ, 08043

Article Info

Received: May 22, 2022
Accepted: June 06, 2022
Published: June 08, 2022

***Corresponding author:** James Espinosa, Department of Emergency Medicine, Rowan University SOM/Jefferson - Stratford, NJ.

Citation: Espinosa J, Lucerna A, Hertz R (2022) Case Report: Cold Thermal Injury and Syncope from Pressurized Dust Cleaning Spray, *Clinical Case Reports and Clinical Study*, 3(7); DOI: <http://doi.org/07.2022/1.138>.

Copyright: © 2022 James Espinosa. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly Cited.

Abstract

Here we present the case of a 21-year-old patient who huffed pressurized dust cleaning spray with subsequent syncope and cold thermal injury to her hands. Air spray cleaners contain halogenated gas, which serves as a propellant to blow dust from electrical and computer devices. The halogenated compound can cause euphoria and can be used as an inhaled abused substance. Use of such substances has been associated with syncope, sudden death, hypoxia and cold thermal injuries. Airway swelling has been reported.

Introduction

Air spray cleaners contain halogenated gas, which serves as a propellant to blow dust from electrical and computer devices. The halogenated compound can cause euphoria and can be used as an inhaled abused substance. The results can include significant morbidity and even mortality. Two associated medical problems are discussed in this case—syncope and cold thermal injury.

The Case:

A 21-year-old female was brought to the emergency department (ED) by emergency medical services (EMS), having been found in a public restroom with a decreased level of consciousness. Two cans of pressurized computer cleaning spray were found with a paper bag and some cloth handtowels. The patient became alert in the ED. She reported huffing the computer cleaning spray and recalled at least two episodes of syncope. Examination of the can showed that the dusting spray contained difluoroethane. She complained of bilateral hand pain. She denied any other complaints. She was not taking any medications. She denied suicidal ideation. She admitted to a two-year history of huffing of computer cleaning spray, sometimes from a paper bag and sometimes through direct inhalation. Her vital signs were within normal limits. On physical exam, deep second-degree burns were noted on both palms. Her physical exam was otherwise within normal limits. Basic laboratory studies were within normal limits. Her ECG showed a sinus rhythm with no acute ST-T abnormalities. After consultation with a burn center, the patient was transferred for further management.

Discussion:

Air spray cleaners contain halogenated gas, which serves as a propellant to blow dust from electrical and computer devices¹. The halogenated compound can cause euphoria and can be used as an inhaled abused substance. Such substances can be used by direct inhalation (sniffing) or through inhalation from a saturated cloth or paper bag--(huffing/bagging).

Inhalation (huffing) of air duster can occur in any age group. It is more common in teenagers and adolescents²⁻⁴. The propellant in air duster is generally 1,1-Difluoroethane (DFE). Some dusting sprays contain trifluoroethane. This compound has been associated with syncope, arrhythmias and even sudden death^{3,5}. The mechanism of an arrhythmia has been shown in animal studies to be sensitization of



the myocardium to catecholamines, probably due to the halogenated hydrocarbon moiety of DFE^{5,6}. Inhalation abuse can cause angioedema³. If a tightly fitting bag is placed over the head with huffing, anoxia can result⁵.

Cold thermal injury has been described in association with DFE containing inhalants¹. The rapid release of pressurized gas causes a cooling effect on the can. Orofacial and digital frostbite has been described as a clue to the detection of the use of such inhalants⁷. Sweating is associated with hydrocarbon use has been hypothesized to allow deeper penetration of the cold injury to the skin⁸.

Inhalants, such as 1,1-Difluoroethane containing products, are considered to be addictive. They can cause a sense of euphoria^{4,5}. They are inexpensive and readily available. Volatile substance abuse is noted to be most common in males between the ages of 14 and 22 years of age^{7,9}. Inhaled hydrocarbons are absorbed through the lungs and exert are readily absorbed by lipids in the brain⁶. Inhalants can affect multiple central neurotransmitters².

In summary, air spray cleaners contain halogenated gas, which serves as a propellant to blow dust from electrical and computer devices. The halogenated compound can cause euphoria and can be used as an inhaled abused substance. The rapid release of pressurized gas causes a cooling effect on the can. Orofacial and digital frostbite has been described as a clue to the detection of the use of such inhalants.

Conflict of Interest: There was no funding related to this case report. The authors declare that they have no conflicts of interest.

References:

1. Avella J, Wilson JC, Lehrer M. Fatal cardiac arrhythmia after repeated exposure to 1,1-difluoroethane (DFE). *Am J Forensic Med Pathol.* 2006 Mar;27(1):58-60.
2. Bonamonte D, Profeta G, Conserva A, Mazzoccoli S, Foti C, Angelini G. Cold burn from contact with a propane and butane gas blend inside a spray canister used as a hooter. *Contact Dermatitis.* 2008 Jul;59(1):61-2. doi: 10.1111/j.1600-0536.2008.01331.x
3. Duncan JR, Lawrence AJ. Conventional concepts and new perspectives for understanding the addictive properties of inhalants. *J Pharmacol Sci.* 2013;122(4):237-43. Epub 2013 Jul 25.
4. Koehler MM, Henninger CA. Orofacial and digital frostbite caused by inhalant abuse. *Cutis.* 2014 May;93(5):256-60.
5. Sakai K, Maruyama-Maebashi K, Takatsu A, Fukui K, Nagai T, Aoyagi M, Ochiai E, Iwadata K. Sudden death involving inhalation of 1,1-difluoroethane (HFC-152a) with spray cleaner: three case reports. *Forensic Sci Int.* 2011 Mar 20;206(1-3):e58-61. doi: 10.1016/j.forsciint.2010.08.026. Epub 2010 Sep 26.
6. Steffee CH, Davis GJ, Nicol KK. A whiff of death: fatal volatile solvent inhalation abuse.
 - a. *South Med J.* 1996 Sep;89(9):879-84.
7. Tormoehlen LM, Tekulve KJ, Nañagas KA. Hydrocarbon toxicity: A review. *Clin Toxicol (Phila).* 2014 Jun;52(5):479-89. doi: 10.3109/15563650.2014.923904.
8. Winston A, Kanzy A, Bachuwa G. Air Duster abuse causing rapid airway compromise. *BMJ Case Rep.* 2015 Jan 7;2015. pii: bcr2014207566. doi: 10.1136/bcr-2014-207566.
9. Xiong Z, Avella J, Wetli CV. Sudden death caused by 1,1-difluoroethane inhalation. *J Forensic Sci.* 2004 May;49(3):627-9.