



X-Linked Retinitis Pigmentosa Revealed by Tapetal Like Reflex: When You All Need for Diagnosis Is a Fundus Picture- A Case Report and Litterature Review

Elkhoyaali Adil

Elkhoyaali.a, military training hospital mohamed v rabat, morocco

Article Info

Received: May 03, 2025

Accepted: May 30, 2025

Published: June 03, 2025

***Corresponding author:** Elkhoyaali Adil, Elkhoyaali.a, military training hospital mohamed v rabat, morocco.

Citation: Elkhoyaali Adil. (2025) "X-Linked Retinitis Pigmentosa Revealed by Tapetal Like Reflex: When You All Need for Diagnosis Is a Fundus Picture- A Case Report and Litterature Review.", *Clinical Medical Case Reports and Case Series*, 2(1); DOI: 10.61148/3065-7644/CMCRCS/012

Copyright: © 2025. Elkhoyaali Adil. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract:

Tapetal-like reflex (TLR) is an unusual, golden, bright, scintillating, particulate reflection on indirect ophthalmoscopy that relatively spares the fovea, similar to that seen in the eyes of some vertebrates.¹ TLR has been described in female carriers of X-linked RP and has also been seen in a healthy young male.²

Keywords: inherited retinal diseases

Introduction:

Retinitis pigmentosa is a group of inherited retinal diseases characterized by degeneration of rod and cone photoreceptors.³ Human inherited retinal degenerations are genetically heterogeneous, with well over 100 genes implicated to date.⁴ Measurements of retinal function, such as the electroretinogram, show that photoreceptor function is usually compromised many years before symptomatic night blindness, visual field scotomas, or decreased visual acuity occur. More than 45 genes have been identified for retinitis pigmentosa.^{5,6}

X-linked Retinitis Pigmentosa GTPase Regulator is a GTPase-binding protein encoded in humans by the RPGR gene, which is located on the X chromosome and is commonly associated with X-linked retinitis pigmentosa (XLRP).⁷ In photoreceptor cells, RPGR is localized in the connecting cilium, which connects the protein-synthesizing inner segment to the photosensitive outer segment and is involved in modulating cargo trafficking between the two segments.⁸

Case Presentation:

We report the case of a 45-year-old woman presented with complaints of decreased vision in the left eye (LE). Her best-corrected visual acuity in the RE was 10/10 and left eye was 06/10. Both eyes (BEs) anterior segment examination was unremarkable. There was no history of night blindness or decreased vision in any of the family members.

Fundus examination revealed presence of retinal pigment epithelial (RPE) hypopigmentation and atrophy at the posterior pole along with an enhanced golden tapetal sheen seen in posterior pole of the left eye (figure 1A).

Fundus autofluorescence revealed a crescent-shaped hyperautofluorescence in LE (figure 1B). Spectral domain optical coherence tomography (SD-OCT) revealed a normal foveal contour in the left eye with thinning of photoreceptor layer to the macula in LE with small drusens (figure 2).

Based on the multimodal imaging findings, we suspected a starting manifestations of retinitis pigmentosa and the patient was advised

for genetic analysis revealing mutation on x-linked RPGR/RP3 GENE then set for observation and a routine follow-up.



Figure1:

A: fundus photo showing tapetal like reflex in left eye

B: fundus autofluorescence showing hyperreflectivity of the posterior pole by atrophy starting of the retinal pigmented epithelium.




Figure 2: macular oct of left eye showing extreme thinning of photoreceptor layer with small drusens

Discussion:

TLR has been described in female carriers of X-linked RP and has also been observed in a healthy young male.³ In addition, abnormal fundus reflections in male patients have been reported in Oguchi disease, X-linked retinoschisis, cone-shaped retinal dystrophy, and early X-linked RP. The TLR has been reported to be located deep to the retinal vasculature and at the level of the outer retina and RPE.³ It may be due to deposits, thickening or degeneration of Bruch's membrane, retinal deposits or an alteration at the level of the RPE-photoreceptor interface.³ We report the presence of an enhanced TLR located in the midperiphery in a patient with sector RP, confirming its association with inherited retinal degenerations; however, the nature and origin of the phenomenon are still not clear and further studies will shed light on this unique finding.

The worldwide prevalence of RP is 1:3000 to 1:7000 including simple, syndromic and systemic disease. There is usually no gender predilection, but males are slightly more affected than females due to the existence of X-linked RP as seen in our patient. RP has no ethnic specificity, but in certain forms related to specific gene

mutations it is more common in conanginous populations (such as the USH3 gene associated with Usher syndrome type III).⁹

Most cases of retinitis pigmentosa are monogenic, but the disease is still very genetically heterogeneous¹⁰. Researchers have identified at least 45 loci at which mutations cause the disease, and these genes together account for the disease in just over half of all patients.¹¹

Since the first report describing linkage of an RP locus to a DNA marker on human chromosome X in 1984¹², more than 40 genes

have been associated with RP. Non-syndromic or "simple" cases may be inherited as autosomal dominant (20-25%), autosomal recessive (15-20%), X-linked recessive (10-15%), or sporadic/simple (30%), and 5% may be early-onset and grouped as part of Leber congenital amaurosis (LCA). Rarer forms also exist, such as X-linked dominant, mitochondrial, and digenic (due to mutations in two different genes).⁹

XLRP is due to mutation of 6 genes located in chromosome X, but only two have been identified so far: the retinitis pigmentosa gtpase

regulator RPGR and the Rp-2 protein RP2.¹³ The mutations in these 2 genes are responsible for 80/100 of clinical cases of XLRP which make them a good target to small molecules drug and gene therapy approaches.¹⁴

Conclusion:

Tapetal like reflex is frequently an incidental fundoscopic finding associated with many retinal diseases like XLRP and may appear years before symptoms and vision loss.

Well understanding of the phenotypes and genes included in RP is the key to be more efficient in developing new treatments of the disease including gene specific approaches, transplantation replacing retinal loss tissue and implanting electrical devices.

References:

1. Kalitzeos A, Samra R, Kasilian M, et al. 2019;39:570–80. Cellular imaging of the tapetal-like reflex in carriers of RPGR-associated retinopathy. *Retina*
2. Schatz P, Bregnhøj J, Arvidsson H, et al. 2012;18:1147–55A tapetal-like fundus reflex in a healthy male: evidence against a role in the pathophysiology of retinal degeneration? *Mol Vis*
3. Meindl A, Dry K, Herrmann K, Manson F, Ciccodicola A, Edgar A, Carvalho MR, Achatz H, Hellebrand H, Lennon A, Migliaccio C, Porter K, Zrenner E, Bird A, Jay M, Lorenz B, Wittwer B, D'Urso M, Meitinger T, Wright A (May 1996). "A gene (RPGR) with homology to the RCC1 guanine nucleotide exchange factor is mutated in X-linked retinitis pigmentosa (RP3)". *Nature Genetics*. 13 (1): 35–42. doi:10.1038/ng0596-35. PMID 8673101. S2CID 31695757.
4. Roepman R, van Duijnhoven G, Rosenberg T, Pinckers AJ, Bleeker-Wagemakers LM, Bergen AA, Post J, Beck A, Reinhardt R, Ropers HH, Cremers FP, Berger W (Jul 1996). "Positional cloning of the gene for X-linked retinitis pigmentosa 3: homology with the guanine-nucleotide-exchange factor RCC1". *Human Molecular Genetics*. 5 (7): 1035–41. doi:10.1093/hmg/5.7.1035. hdl:2066/22748. PMID 8817343.
5. Murga-Zamalloa CA, Atkins SJ, Peranen J, Swaroop A, Khanna H (Sep 2010). "Interaction of retinitis pigmentosa GTPase regulator (RPGR) with RAB8A GTPase: implications for cilia dysfunction and photoreceptor degeneration". *Human Molecular Genetics*. 19 (18): 3591–8. doi:10.1093/hmg/ddq275. PMC 2928130. PMID 20631154. Jump up to: a b "Entrez Gene: RPGR retinitis pigmentosa GTPase regulator".
6. Khanna H (October 2015). "Photoreceptor Sensory Cilium: Traversing the Ciliary Gate". *Cells*. 4 (4): 674–86. doi:10.3390/cells4040674. PMC 4695852. PMID 26501325.
7. Churchill JD, Bowne SJ, Sullivan LS, Lewis RA, Wheaton DK, Birch DG, et al. (February 2013). "Mutations in the X-linked retinitis pigmentosa genes RPGR and RP2 found in 8.5% of families with a provisional diagnosis of autosomal dominant retinitis pigmentosa". *Investigative Ophthalmology & Visual Science*. 54 (2): 1411–6. doi:10.1167/iovs.12-11541. PMC 3597192. PMID 23372056.
8. Glomset JA, Farnsworth CC (1994). "Role of protein modification reactions in programming interactions between ras-related GTPases and cell membranes". *Annual Review of Cell Biology*. 10: 181–205. doi:10.1146/annurev.cb.10.110194.001145. PMID 7888176.
9. Stefano Ferrari*,¹, Enzo Di Iorio¹, Vanessa Barbaro¹, Diego Ponzin¹, Francesco S. Sorrentino² and Francesco Parmeggiani²
10. Hartong, D. T., Berson, E. L., & Dryja, T. P. (2006). Retinitis pigmentosa. *The Lancet*, 368(9549), 1795–1809. doi:10.1016/s0140-6736(06)69740-7
11. Hardcastle AJ, Thiselton DL, Zito I, et al. Evidence for a new locus for X-linked retinitis pigmentosa (RP23). 2000; 41: 2080–86. *Invest in Ophthalmol Vis Sci*
12. Gorbatuyk, 2010M.S, Knox.T, La Vail,M.M, Gorbatuyk,O.S ;Lewin,A.S, Restauration of visual fonction in P23H rhodopsin transgenic rats by gene delivery of bip/grp78, *proc.natl. acad.sci USA*
13. Vettel.S, Wittinghoffer, 2009,13(10),1239-1251A.RPGR and RP2: targets for the treatment of x-linked Retinitis pigmentosa? *Expert.Opin.Ther.Targets*
14. Vervoort.R; Lenon, 2000,25(4),462-466 A. ;Bird,A.C ;Tulloch,B ;Axton,R ;Miano,M.G,Mututinal hospot within a new RPGR exon in X-linked retinitis pigmentosa m.Nat,Genet.